

Possible Evidence for a Chiral Axial-Vector State in the D Meson System

Kenji YAMADA*, Muneyuki ISHIDA[¶], Shin ISHIDA[†], Daiki ITO[†],
Toshihiko KOMADA* and Hiroshi TONOOKA[†]

**Department of Engineering Science, Junior College Funabashi Campus
Nihon University, Funabashi 274-8501, Japan*

¶Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan

*†Atomic Energy Research Institute, College of Science and Technology
Nihon University, Tokyo 101-8308, Japan*

We reanalyze the $D^{*+}\pi^-$ mass spectrum from CLEO II by the VMW method in order to examine the existence of a chiral axial-vector state, which is predicted in a covariant level-classification scheme recently proposed, other than normal orbitally-excited P -wave states in the D meson system. A result of the present analysis seems to suggest that there exists an extra axial-vector meson, in addition to the two normal ones, in a similar mass region.

§1. Introduction

In the constituent quark model, together with heavy quark symmetry, the lowest-lying positive parity excitations of heavy-light $Q\bar{q}$ meson systems are expected, in the limit $m_Q \rightarrow \infty$, to be two degenerate spin doublets with the total angular momentum $j_q = 1/2$ and $3/2$ of the light quark, that is, four orbitally-excited states with $L = 1$ labeled as

$$\begin{aligned} j_q L_J = {}^{1/2}P_0, {}^{1/2}P_1 & \quad \text{for the } j_q = \frac{1}{2} \text{ doublet,} \\ = {}^{3/2}P_1, {}^{3/2}P_2 & \quad \text{for the } j_q = \frac{3}{2} \text{ doublet.} \end{aligned}$$

In this limit heavy quark symmetry further requires that the $j_q = 1/2$ states decay to ${}^{1/2}S_0 + \pi$ or ${}^{1/2}S_1 + \pi$ only in an S -wave, while the $j_q = 3/2$ states decay only in a D -wave. It is therefore expected that the decay widths of the $j_q = 1/2$ and $3/2$ states are broad and narrow, respectively.

On the one hand, a covariant level-classification scheme of quark-antiquark meson systems has been proposed, which gives them a covariant quark representation with definite Lorentz and chiral transformation properties.¹⁾ In this scheme, assuming that chiral symmetry for the light-quark component in heavy-light meson systems is effective, the existence of extra scalar and axial-vector states is predicted, respectively, as chiral partners of the ground-state pseudoscalar and vector mesons. These what we call chiral scalar and axial-vector mesons are distinguished from the above-mentioned P -wave states, since the chiral scalar state is an analogue of the $\sigma(400 - 600)$ meson, which is difficult to be interpreted as the 3P_0 state, as a chiral partner of the π meson in the light-quark system.

In this report we present a possible evidence for the chiral axial-vector state, D_1^χ , in the D meson system.

§2. Reanalysis of the $D^{*+}\pi^-$ mass spectrum from CLEO II

We reanalyze the $D^{*+}\pi^-$ mass spectrum, published by CLEO Collaboration,²⁾ by the VMW method in which the production amplitude is expressed by a sum of Breit-Wigner amplitudes for relevant resonances.

In the present analysis we take into account the four states $D_2^*(3/2 P_2)$, $D_1(3/2 P_1)$, $D_1^*(1/2 P_1)$ and D_1^χ which can decay to $D^*\pi$. Then, following the VMW method, the production amplitude is given by

$$|A(s)|^2 = \left| r_1 e^{i\theta_1} \Delta_{D_1^\chi}(s) + r_2 e^{i\theta_2} \Delta_{D_1^*}(s) \right|^2 + \left| r_3 e^{i\theta_3} \Delta_{D_1}(s) \right|^2 + \left| r_4 e^{i\theta_4} \Delta_{D_2^*}(s) \right|^2,$$

$$\Delta_R(s) = \frac{-m_R \Gamma_R}{s - m_R^2 + i m_R \Gamma_R},$$

where $r_1, \dots, r_4$ and $\theta_1, \dots, \theta_4$ are the production couplings and phases of respective resonances, and we assumed that D_1^χ and D_1^* decay only through an S -wave, while D_1 only through a D -wave. The background $D^*\pi$ mass distribution is fit with a five-parameter threshold function given by

$$BG = \alpha (\Delta M)^\beta \exp \left[-\gamma_1 \Delta M - \gamma_2 (\Delta M)^2 - \gamma_3 (\Delta M)^3 \right],$$

$$\Delta M = M(D^*\pi) - m_{D^*} - m_\pi,$$

where the parameters α , β , γ_1 , γ_2 and γ_3 are fixed through the fit to the total $D^*\pi$ mass spectrum.

Using the above production amplitude and background, we fit the $D^*\pi$ mass spectrum in the following three cases:

- (a) High-mass D_1^* with a mass of $2500 < m_{D_1^*} < 2600$ in MeV,
- (b) Low-mass D_1^* with a mass of $2350 < m_{D_1^*} < 2500$ in MeV,
- (c) No D_1^χ and D_1^* .

Here the case (c) corresponds to the original analysis by CLEO Collaboration, though the background parametrization is somewhat different. The results of fits are shown in Fig. 1 and obtained values of the resonance parameters are given in Table I. In both the fits with high- and low-mass D_1^* , it is found that the mass and width of D_1^χ are ≈ 2310 MeV and ≈ 20 MeV, respectively, and those of D_1 and D_2^* are similar to the values reported so far. For the mass and width of D_1^* we obtain ≈ 2600 MeV and ≈ 200 MeV in the high-mass fit, while ≈ 2420 MeV and ≈ 200 MeV in the low-mass fit.

In all the three cases of fits the χ^2/N_{dof} is best for the high-mass D_1^* fit, though they are not so different from each other. It would be worth while noting that the two cases of fits with D_1^χ seem to describe the data better than the fit without D_1^χ in the mass region 2.15–2.5 GeV, where there appears to be an excess of data events at the mass 2.31–2.33 GeV.

§3. Theoretical remarks on the results

We consider the mass splitting and mixing of P -wave meson multiplets, based on the Breit-Fermi Hamiltonian with vector-gluon and long-range-scalar exchange,




Fig. 1. The results of the fits to the $D^{*+}\pi^-$ mass spectrum with (a) high-mass D_1^* , (b) low-mass D_1^* , and (c) no D_1^X and D_1^* .

Table I. Values of the resonance parameters and χ^2/N_{dof} obtained from the respective fits.

	(a) Fit with high-mass D_1^*		(b) Fit with low-mass D_1^*		(c) Fit without D_1^X and D_1^*	
State	Mass (MeV)	Width (MeV)	Mass (MeV)	Width (MeV)	Mass (MeV)	Width (MeV)
D_1^X	2308	18.7	2307	17.4	—	—
D_1^*	2596	199	2421	199	—	—
D_1	2421	34.5	2421	27.0	2421	27.5
D_2^*	2472	35.0	2468	35.0	2466	35.0
χ^2/N_{dof}	57.7/52		58.0/52		66.2/59	

where we ignore the P -wave D_1 and D_1^* states mixing with the chiral D_1^X state. Taking a static potential due to single vector-gluon exchange to be $-4\alpha_s/3r$, the spin-dependent part of the Hamiltonian for P -wave states can be expressed, to first order $1/m_Q$, as

$$\delta\mathcal{H} = C_q \mathbf{L} \cdot \mathbf{S}_q + C_Q (\mathbf{L} \cdot \mathbf{S}_Q + S_T), \quad S_T = 3(\mathbf{S}_q \cdot \hat{\mathbf{r}})(\mathbf{S}_Q \cdot \hat{\mathbf{r}}) - \mathbf{S}_q \cdot \mathbf{S}_Q$$

with

$$C_q = \left(\frac{1}{2m_q^2} + \frac{1}{m_q m_Q} \right) \left\langle \frac{4\alpha_s}{3r^3} \right\rangle - \frac{1}{2m_q^2} \left\langle \frac{1}{r} \frac{dV_s}{dr} \right\rangle,$$

$$C_Q = \frac{1}{m_q m_Q} \left\langle \frac{4\alpha_s}{3r^3} \right\rangle,$$

where $V_s(r)$ is the static potential due to long-range-scalar exchange and the spin-spin interaction is neglected because of its contact nature. The Hamiltonian $\delta\mathcal{H}$ gives rise to the mass splittings among P -wave multiplets and the mixing between the $^{3/2}P_1$ and $^{1/2}P_1$ states. Treating $\delta\mathcal{H}$ as a first-order perturbation and using the mass values of D_2^* , D_1 and D_1^* obtained in the high-mass D_1^* fit, we find ≈ 2470 MeV for the mass of $D_0^*(^{1/2}P_0)$ and $\phi - \phi_{HQ} \approx -3.64^\circ$ for the deviation of the $D_1(^{3/2}P_1) - D_1^*(^{1/2}P_1)$ mixing angle from the heavy-quark-symmetry limit with the parameter values of the unperturbed mass $M_0 = 2490$ MeV common to all four states, $C_q = -73.81$ MeV and $C_Q = 47.15$ MeV, where we have chosen a solution with $C_Q > 0$ in accord with the above definition of C_Q . For the low-mass D_1^* case of fits there is no solution.

§4. Concluding remarks

We have reanalyzed the $D^{*+}\pi^-$ mass spectrum published by CLEO Collaboration and found a possible evidence for the chiral axial-vector meson D_1^χ with a mass and width of ≈ 2310 MeV and ≈ 20 MeV, respectively. We have also found the mass and width of D_1^* to be ≈ 2600 MeV and ≈ 200 MeV, together with similar masses and widths of D_1 and D_2^* to those reported so far, among which the spin-dependent Hamiltonian arising from one-vector-gluon and long-range-scalar exchange could account for the mass splittings. To confirm the existence of D_1^χ it goes without saying that further analyses, including other experimental data with high statistics, are necessary in a more precise way.

Furthermore, in establishing the covariant level-classification scheme of meson systems it is important to examine the existence of the chiral scalar meson D_0^χ as well as B_0^χ and B_1^χ in the B meson system. An analysis of the $B\pi$ mass spectrum, to study the existence of B_0^χ , is in progress and its preliminary result has been presented.³⁾

References

- [1] Ishida, S., Ishida, M., and Maeda, T., Prog. Theor. Phys. **104**, 785-807 (2000).
- [2] CLEO Collaboration, Phys. Letters B **331**, 236-244 (1994); **342**, 453(E) (1995).
- [3] Ishida, M., and Ishida, S., these proceedings.